

# Practical BSE Calculations with BerkeleyGW + Octopus

---



Felipe H. da Jornada

Department of Physics – University of California, Berkeley  
and Lawrence Berkeley National Laboratory



David A. Strubbe

Department of Material Science and Engineering –  
Massachusetts Institute of Technology

TDDFT Benasque Workshop – Jan. 12<sup>th</sup>, 2014

# Summary


---

#1 – Theory and Algorithms

#2 – Typical BSE Workflow in BerkeleyGW


#3 – Issues Unique to the BSE Code

# Theory and Algorithms



# Theory Review: Optical Absorption


## No electron-hole interactions



$$H_{int} \sim \mathbf{A} \cdot \hat{\mathbf{v}}$$

$$\epsilon_2(-\mathbf{q}, \omega) \propto \sum_{v\mathbf{c}\mathbf{k}} |\langle v\mathbf{k} + \mathbf{q} | \hat{\mathbf{v}} | c\mathbf{k} \rangle|^2 \delta[\omega - (E_{c\mathbf{k}} - E_{v\mathbf{k}+\mathbf{q}})]$$

## With electron-hole interactions



$$\epsilon_2(-\mathbf{q}, \omega) \propto \sum_S |\langle 0 | \hat{\mathbf{v}} | S \rangle|^2 \delta[\omega - \Omega_S]$$

$$|S\rangle = \sum_{v\mathbf{c}\mathbf{k}} A_{v\mathbf{c}\mathbf{k}}^S |v\mathbf{k} + \mathbf{q}\rangle \otimes |c\mathbf{k}\rangle$$

Solutions of the Bethe-Salpeter equation (BSE)

# Bethe Salpeter Equation (BSE)

- ▶ From Prof. Tokalty's talk to BerkeleyGW:

$$L = (1 - L_0 K)^{-1} L_0$$

$$L^{-1} = L_0^{-1} - K$$



Spectral Representation:

$$-L_0^{-1} = E_c - E_v - \omega$$

$$-L^{-1} = \boxed{E_c - E_v + K} - \omega$$

- ▶ Absorption spectrum with excitonic effects → diagonalize BSE Hamiltonian:

$$[H]_{(vc\mathbf{k}), (v'c'\mathbf{k}' )}$$

$$[H] = [E_c - E_v] + [K]$$

↖ diagonal  
~ kinetic term

← dense "kernel"  
~ potential term

Challenge: compute **quasiparticle corrections** and **kernel** matrix elements on a very fine k-grid!

# BerkeleyGW Interpolation Scheme

BerkeleyGW solution:  
Interpolate QP energies and BSE kernel

- ▶ Step 1: Expand fine WFNs in terms of coarse WFNs


$$u_{n\mathbf{k}_{\text{fi}}} = \sum_{n'} C_{n,n'}^{\mathbf{k}_{\text{co}}} u_{n'\mathbf{k}_{\text{co}}}$$

- ▶ Step 2: Interpolate QP energies and matrix elements

$$\langle \underline{v\mathbf{k}_{\text{fi}}} | K | \underline{v'\mathbf{k}'_{\text{fi}}} \rangle = \sum_{n_1, n_2, n_3, n_4} C_{c, n_1}^{\mathbf{k}_{\text{co}}} C_{v, n_2}^{*\mathbf{k}_{\text{co}}} C_{c', n_3}^{*\mathbf{k}'_{\text{co}}} C_{v', n_4}^{\mathbf{k}'_{\text{co}}} \langle n_2 n_1 \underline{\mathbf{k}_{\text{co}}} | K | n_4 n_3 \underline{\mathbf{k}'_{\text{co}}} \rangle$$

# BerkeleyGW Interpolation Scheme


- ▶ In practice: trading bands for k-points



- ▶ How to get a good interpolation?
  - ▶ Include a large number of bands from the coarse grid!

# BerkeleyGW QP Interpolation

- ▶ BerkeleyGW also performs a linear interpolation for QP corrections.
- ▶ Linear interpolation + expansion over bands:
  - ▶ Captures (nk)-dependent QP correction and band crossing
  - ▶ Very smooth interpolation of band structure
  - ▶ Robust scheme, and very few parameters
- ▶ This is how `inteqp.x` works!



D. Qiu, F. H. da Jornada, S. G. Louie, PRL **111**, 216805 (2013)




# Theory and Algorithms

## Key Points

- Interpolation scheme: trading bands for k-points
- Interpolation of kernel and QP corrections

# Typical BSE Workflow in BerkeleyGW



# Goal

---

Goal: Diagonalize BSE Hamiltonian on a fine grid

$$[H]_{fi} = [E_c - E_v]_{fi} + [K]_{fi}$$

# BerkeleyGW Workflow

---

sigma.x

**Step 1:** Calculate QP-corrected band structure on a coarse grid

$$\{E_c\}_{co}, \{E_v\}_{co},$$

kernel.x

**Step 2:** Calculate BSE kernel on the same coarse grid

$$[K]_{co}$$

absorption.x

**Step 3:** Interpolate to a fine k-grid and build BSE Hamiltonian...

$$[H]_{co} \Rightarrow [H]_{fi}$$

... and diagonalize BSE Hamiltonian

$$\text{evals } [H]_{fi} \Rightarrow \varepsilon_2$$

(Not shown: mean-field, epsilon, convergence)

# I. Sigma

---

sigma.x

**Step I:** Calculate QP-corrected band structure on a coarse grid

$$\{E_c\}_{co}, \{E_v\}_{co},$$

- ▶ Same procedure done in previous sessions.
- ▶ Recommended: eqp.dat
  - ▶ Calculate QP energies on all k-points from WFN inner.
  - ▶ Use the script eqp.py to generate eqp.dat file → no human intervention!
- ▶ Also possible: scissors operators, less reliable (not covered here)

# I. Sigma

Sample sigma.inp (assuming we are using eqp.dat)

```
screened_coulomb_cutoff <?>
bare_coulomb_cutoff <?>

number_bands <?>
band_occupation <?>

band_index_min <?>
band_index_max <?>


screening_semiconductor
number_kpoints <?>
begin kpoints
  <put all k-points from WFN_INNER here>
end
```

Note the two  
different # of bands:

$$E_{n\mathbf{k}}^{\text{QP}} = \langle n\mathbf{k} | \Sigma | n\mathbf{k} \rangle$$



Remember to  
calculate Sigma on  
more bands because  
of the interpolation!




## 2. Kernel

kernel.x

**Step 2:** Calculate BSE kernel on the same coarse grid

$$[K]_{co}$$

- ▶ Time consuming!
  - ▶ Computes  $(n_v n_c n_k)^2$  matrix elements



- ▶ Recommended:
  - ▶ Use same WFN co as in Sigma (WFN inner)

## 2. Kernel

---

Sample kernel.inp

```
number_val_bands <?>
number_cond_bands <?>
screened_coulomb_cutoff <?>

<?>_symmetries_coarse_grid
screening_<?>
```

Remember to calculate  
Kernel on more bands  
because of the  
interpolation!

Must be  $\leq$  than the #  
of bands used in Sigma.

You'll typically want to use symmetries  
here, so put:  
use\_symmetries\_coarse\_grid



# 3. Absorption

---

absorption.x

**Step 3:** Interpolate to a fine k-grid and build BSE Hamiltonian...

$$[H]_{co} \Rightarrow [H]_{fi}$$

... and diagonalize BSE Hamiltonian

$$\text{evals } [H]_{fi} \Rightarrow \varepsilon_2$$

- ▶ Absorption needs same coarse WFN\_co from Kernel/Sigma
- ▶ Absorption also need two fine WFN files:
  - ▶ WFN\_fi: for conduction states
  - ▶ WFNq\_fi: for valence states
- ▶ Good practice: use randomly-shifted k-grids
  - ▶ This maximizes the number of inequivalent transitions you capture.

# 3. Absorption

Sample absorption.inp

```
diagonalization
```

```
number_val_bands_coarse <?>
number_cond_bands_coarse <?>
number_val_bands_fine <?>
number_cond_bands_fine <?>
```

```
coarse_grid_points <?>
```


```
use_symmetries_coarse_grid
no_symmetries_fine_grid
no_symmetries_shifted_grid
```

```
screening_semiconductor
```

```
use_velocity
q_shift 0.0 0.0 0.001
```

```
gaussian_broadening
energy_resolution 0.15
```

```
eqp_co_corrections
```



Same as used in kernel.

We interpolate to these bands!

How many k-points in the coarse grid after unfolding BZ?

Typical values.


Recommended!

$$\mathbf{k}_{\text{WFN}_{fi}} + \mathbf{q}_{\text{shift}} = \mathbf{k}_{\text{WFN}_{qfi}}$$


Broaden each delta function.

Interpolate eqp\_co.dat to eqp.dat

# 3. Absorption – Workflow




# Typical BSE Workflow in BerkeleyGW



## Key Points

- BSE codes separated into two parts:
  - Kernel.x: calculates kernel on coarse grid
  - Absorption.x: interpolates and diagonalizes [H]
- `number_*_bands_coarse`, `number_*_bands_fine`.



# Issues Unique to the BSE Code

1. Velocity Operator
2. Finite Systems + Octopus
3. Estimating the Quality of the Interpolation
4. Analyzing Exciton Files
5. Convergence!

# I. Velocity Operator

$$\varepsilon_2(-\mathbf{q}, \omega) \propto \sum_S |\langle 0 | \hat{v} | S \rangle|^2 \delta[\omega - \Omega_S] \quad \langle 0 | \hat{v} | S \rangle = \sum_{v\mathbf{c}\mathbf{k}} A_{v\mathbf{c}\mathbf{k}}^S \langle v\mathbf{k} + \mathbf{q} | \hat{v} | c\mathbf{k} \rangle$$

- ▶ Because of non-local pseudopotential and QP corrections, the velocity operator is not the same as the momentum!

$$\hat{v} = i[H, \hat{r}] = \hat{p} + i[V, \hat{r}]$$

## use\_velocity

- ▶ Recommended option!
- ▶ Needs WFN<sub>fi</sub> and WFN<sub>q\_fi</sub>.
- ▶ Specify q-shift:  $\mathbf{k}_{\text{WFN}_{\text{fi}}} + \mathbf{q}_{\text{shift}} = \mathbf{k}_{\text{WFN}_{\text{qfi}}}$

$$\langle 0 | \hat{v} | S \rangle \approx \frac{\Omega_S}{q} \sum_{v\mathbf{c}\mathbf{k}} A_{v\mathbf{c}\mathbf{k}}^S \langle v\mathbf{k} + \mathbf{q} | e^{-i\mathbf{q}\cdot\mathbf{r}} | c\mathbf{k} \rangle$$

## use\_momentum

- ▶ Not recommended!
- ▶ Needs only WFN.
- ▶ Specify polarization  $\mathbf{e}_\lambda$  of  $\hat{v}$ .

$$\langle v\mathbf{k} + \mathbf{q} | \hat{v} | c\mathbf{k} \rangle \approx \langle v\mathbf{k} | \hat{p}_\lambda | c\mathbf{k} \rangle$$

## 2. Finite Systems + Octopus

- ▶ For finite systems, Octopus can calculate the single-particle velocity matrix elements directly!
  - ▶ No need for two different WFN files and a q-shift in the absorption code.
  - ▶ BerkeleyGW can calculate  $\varepsilon_2(\mathbf{q} = 0)$  “exactly”.
- ▶ In BerkeleyGW:

```
...  
read_vmtxel  
use_momentum  
polarization <...>  
  
skip_interpolation  
eqp_corrections  
...
```

Read velocity matrix elm from file

Tell BerkeleyGW that there's no q-shift.

Note: this doesn't affect calculation of matrix elms because of the read\_vmtxel flag.

There are no k-points, so there's no interpolation! Use directly the eqp.dat file (don't interpolate eqp\_co.dat → eqp.dat)

# 3. Quality of the Interpolation

- ▶ How to measure the quality of WFN expansion?
- ▶ If we include  $\infty$  bands:

$$\sum_{n'} |C_{n,n'}^{\mathbf{k}_{co}}|^2 = 1$$

- ▶ Finite basis set – normalization is reported in files `d?mat_norm.dat`:

```
----- Norm of dvv matrices : Spins = 1 -----
          k-point          ik_co  v  dist  |dvv|^2
-----
( 0.059 , 0.046 , 0.039 )    1    1  0.054  0.987006
( 0.059 , 0.046 , 0.039 )    1    2  0.054  0.953488
( 0.059 , 0.046 , 0.039 )    1    3  0.054  0.892665
( 0.059 , 0.046 , 0.164 )    2    1  0.139  0.923182
```

- ▶ How to get a good interpolation?
  - ▶ Include a large number of bands from the coarse grid!
  - ▶ Start from a fine enough grid

Before renormalization of coefficients.




# 4. Analyzing Excitons

---

- ▶ Optical spectrum  $\varepsilon_2(\omega)$ ,  $\varepsilon_1(\omega)$ :
  - ▶ absorption\_noeh.dat: GW-RPA without local fields
  - ▶ absorption\_eh.dat: GW-BSE with local fields
  
- ▶ Eigenvalues of the BSE equation  $\Omega_S$ :
  - ▶ eigenvalues.dat: useful to see if there are degeneracies, splitting, etc.
  
- ▶ Where the exciton is coming from:
  - ▶ `summarize_eigenvectors.x`
  - ▶ Need to set the flag `write_eigenvalues` in `absorption.inp`

# 5. Convergence!

- ▶ There are 4 convergence parameters in a typical BSE calculation:
  - ▶ # of k-points in the fine grid
  - ▶ # of bands in the fine grid
  - ▶ # of k-points in the coarse grid
  - ▶ # of bands in the coarse grid



D. Qiu, F. H. da Jornada, S. G. Louie, PRL  
111, 216805 (2013).

Make sure you converge your calculations!

# Issues Unique to the BSE Code

## 1. Velocity operator

### Key Points

- Why we need 2 WFN files: velocity operator
- **Convergence!**

## 5. Convergence!

# Let's Put the Donkey to Work!



## BerkeleyGW

*Predicting quasiparticle band structures since 1985.*